

221031 WART CORN AND CALLUS CREAM

What is this preparation?

This preparation is custom-formulated specifically for you to better treat your individual needs. The unique combination of active and inactive ingredients has been selected to produce an outcome that is clinically different to that which is commercially available. For this reason, no standardized information or literature is available for your final preparation. Custom-created preparations allow us to meet your individual needs without being restricted to off-the-shelf medicines and devices.

What is this preparation for?

This is a custom-formulated preparation designed to better meet your individual needs. This practice's desire is to help you achieve better outcomes. Its active ingredients will work together toward this goal.

Cimetidine is an FDA approved anti-acid medicine, but several recent studies showed that its off label use topically in conjunction with other ingredients may be beneficial for the treatment of warts.

Lidocaine is a commercially available drug in several topical forms and strengths used as local anesthetic.

Salicylic Acid is a key ingredient in many skin-care and dermatology products for the treatment of seborrheic dermatitis, acne, psoriasis, keratosis pilaris, acanthosis nigricans, ichthyosis, warts, etc. It is used in various concentrations depending on the purpose. Being in the beta hydroxy acid category, Salicylic Acid acts as a keratolytic, exfoliating or peeling agent.

How to use this preparation?

Use this preparation exactly as your doctor instructed you.

Additional directions:

How to store this preparation

This preparation should be stored in a cool, dry and dark place.

Warning: You may have an allergic reaction to any of the ingredients in this custom-formulated preparation. It is advisable that you first test the preparation before applying it on a large area of the body. Apply a very small amount of cream/lotion/gel/solution on a quarter size area on your forearm to test. Let it stay for about 30 minutes. If you have a fast reaction type of allergy, the area will be red and itchy. If you have a slow reaction type of allergy, the same symptoms may occur after 48 hours. If any of these situations occur, stop using the preparation and call your health care practitioner. The absence of any reaction to this test is no guarantee that an allergic reaction may not occur during future uses of this preparation.